

# Dark matter dynamics in Galactic center

Eugene Vasiliev and Maxim Zelnikov  
Lebedev Physical Institute, Moscow

Nbody 2008 Workshop, August 2008 Turku, Finland

# Why the dark matter life and adventures in Galactic center are interesting?


- There is much DM in the center:  $\rho(r) \sim r^{-1}$  originally or even steeper after baryonic compression
- High density implies high annihilation rate:  $J \sim \int \rho^2(r) d^3r$
- The distribution of DM changes greatly during galactic lifetime because of gravitational interaction with stars
- The supermassive black hole in galactic center is nourished by DM as well as by ordinary matter

# What processes are significant for dark matter evolution in Galactic center?

- Baryonic compression

For circular orbits the angular momentum is conserved:


$$L^2 = v_{circ}^2 r^2 = M_{dm,in}(r) r \Rightarrow (M_{dm,fin}(r') + M_{bar}(r')) r'$$



In general case, the radial action is also adiabatically conserved.

# What processes are significant for dark matter evolution in Galactic center?

- Dark matter annihilation near a black hole  
DM density drops at small radii




In the case of circular orbits,  
the annihilation plateau is formed

$$\rho_a = \frac{m_\chi}{\langle\sigma v\rangle t}$$

In more realistic cases (e.g.  
isotropic velocity distribution)  
a weak cusp is formed

# What processes are significant for dark matter evolution in Galactic center?

- Dark matter scattering by stars and absorption by SMBH



Merritt, Harfst & Bertone 2007

Bahcall & Wolf 1976

Lightman & Shapiro 1977

Cohn & Kulsrud 1978

} stars

Merritt 2004

Ilyin, Zybin & Gurevich 2004

Bertone & Merritt 2005

Vasiliev & Zelnikov 2008

} dark matter

# Fokker-Planck treatment of DM evolution

## Assumptions:

- spherical symmetry:  
using variables  $E$  (energy) and  $L$  (angular momentum)
- single supermassive black hole in the galactic center
- orbit-averaged kinetic equation  
(distribution function changes slowly compared to dynamical time)

## Additionally

evolution considered both within and outside SMBH radius of influence ( $r_h = 2$  pc for Milky Way);

relaxation timescale  $t_r \sim 2.5$  Gyr within  $r_h$  and quickly grows outside  $r_h$

# Fokker-Planck treatment of DM evolution

The kinetic equation:

$$\frac{\partial f(E, R, t)}{\partial t} = \mathcal{G}^{-1} \frac{\partial}{\partial \xi_\alpha} \left\{ \mathcal{G} \left[ D_{\alpha\beta} \frac{\partial f}{\partial \xi_\beta} - D_\alpha f \right] \right\} - S_{ann}[f], \quad \mathcal{G} - \text{jacobian},$$

$\xi_\alpha = \{E, R\}$  are phase space variables:  $E$  is energy and  $R = \frac{L^2}{L_c(E)^2} \in [0..1]$  is scaled angular momentum squared ( $L_c(E)$  is maximal angular momentum for given energy  $E$ ).

$D_\alpha$  и  $D_{\alpha\beta}$  are drift and diffusion coefficients due to scattering on stars.

$D_\alpha \propto \frac{m}{m_\star} \Rightarrow$  drift is insignificant for dark matter.

$S_{ann}$  is orbit-averaged annihilation loss term.

# Fokker-Planck treatment of DM evolution

The black hole imposes boundary condition at  $R=R_{min}(E)$  :

$$\left( f - \alpha R_{min} \frac{\partial f}{\partial R} \right) \Big|_{R=R_{min}} = 0 .$$

$$\alpha \approx \begin{cases} \sqrt{q} & , q < 1 ; \\ q & , q > 1 ; \end{cases}$$

$$q = (D_{RR}/R) \Big|_{R \rightarrow 0} T_{orb}(E) / R_{min}(E)$$


Low  $E$ :  $q \ll 1$  (empty loss cone)

High  $E$ :  $q \gg 1$  (full loss cone)

The distribution function at fixed  $E$  and small  $R$  is close to logarithm:

$$f(R) = f_{min} \cdot \left( 1 + \frac{1}{\alpha} \ln \frac{R}{R_{min}} \right).$$

↑  
(capture boundary)



For stars this allows to reduce the equation to 1-dimensional (for energy only)


# Fokker-Planck treatment of DM evolution

## Main features of our consideration:

- two-dimensional (E and R):  
diffusion coefficients vary significantly with R, so R-averaging and reduction to one-dimensional equation for energy is questionable
- broad class of initial DM profiles (NFW, Moore, etc.)  
allows comparison of different models and detects common factors
- analytic approximations for early ( $t \ll t_r$ ) and late ( $t \gg t_r$ ) times;  
analytic consideration of 1-dim. simplified equations for E and R;  
full 2-dim. equation is solved numerically.


Main difference from stars: no (quasi)-stationary DM profile (as opposed to Bahcall-Wolf cusp) because of constant heating of DM particles by stars [energy equipartition of light (DM) and heavy (star) species]

# Dark matter density evolution



example for initial DM profile of NFW ( $\rho \sim r^{-1}$ )

# Dark matter budget after $10^{10}$ yr of evolution



example for initial DM profile of NFW ( $\rho \sim r^{-1}$ )

# Conclusions

- Dark matter distribution in the Galactic center significantly evolved since Galaxy formation; the evolution is due to gravitational scattering on stars, capture by black hole and annihilation
- Evolution is followed by method of Fokker-Planck equation in two-dimensional phase space of energy and angular momentum
- Differences between various initial DM density profiles are greatly reduced after several relaxation times
- DM density determines intensity of annihilation radiation from the Galactic center; predictions from the calculations are consistent with observational constraints

\*\*\*